

# Multi-periodic climate dynamics: spectral analysis of long-term instrumental and proxy temperature records

geschrieben von H.-j. Lüdecke, A. Hempelmann, C.o. Weiss | 2. November  
2013

H.-J. Lüdecke, A. Hempelmann, and C.O. Weiss: Multi-periodic climate  
dynamics: spectral analysis of long-term instrumental and proxy  
temperature records, *Clim. Past.*, 9, 447-452, 2013,  
doi:10.5194/cp-9-447-2013, [www.clim-past.net/9/447/2013/](http://www.clim-past.net/9/447/2013/)

## Abstract:

The longest six instrumental temperature records of monthly means reach back maximally to 1757 AD and were recorded in Europe. All six show a V-shape, with temperature drop in the 19th and rise in the 20th century. Proxy temperature time series of Antarctic ice cores show this same characteristic shape, indicating this pattern as a global phenomenon. We used the mean of the 6 instrumental records for analysis by discrete Fourier transformation (DFT), wavelets, and the detrended fluctuation method (DFA). For comparison, a stalagmite record was also analyzed by DFT. The harmonic decomposition of the above mentioned mean shows only 6 significant frequencies above periods over 30 yr. The Pearson correlation between the mean, smoothed by a 15 yr running average (boxcar) and the reconstruction using the 6 significant frequencies yields  $r = 0.961$ . This good agreement has a 99.9% confidence level confirmed by Monte Carlo simulations. It shows that the climate dynamics is governed at present by periodic oscillations. We find indications that observed periodicities result from intrinsic dynamics.